

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



QuickBayt[®]

FLY BAIT

Active Constituents: 5.0 g/kg IMIDACLOPRID
1.0 g/kg (Z)-9-TRICOSENE

GROUP 4A INSECTICIDE

Ready-to-use fly bait granules for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTION FOR USE

QuickBayt contains BITREX, an acutely bitter agent which reduces the possibility of accidental ingestion by children, domestic animals or wildlife.

GENERAL INSTRUCTIONS

QuickBayt can be applied in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. For best results time applications to begin at the start of the season before fly populations have reached their peak.

Fly control should be considered part of an integrated strategy which incorporates good sanitation, appropriate management of breeding sites (eg. manure and other decaying organic matter) and exclusion measures for preventing access of flies into internal areas.

NOTE: This product is not attractive to all species of flies and control of non-target species may require other means of control.

Insecticide Resistance Warning

For insecticide resistance management, QuickBayt Fly Bait is a Group 4A insecticide.

Some naturally occurring insect biotypes resistant to QuickBayt and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if QuickBayt or other Group 4A insecticides are used repeatedly. The effectiveness of QuickBayt on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of QuickBayt to control resistant insects.

For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science or Bayer Animal Health representative.

PRECAUTIONS

Do NOT allow contact with raw or processed food, utensils or packaging.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Do NOT place baits in locations accessible to domestic animals, livestock or birds. Do NOT contaminate streams, rivers or waterways with the chemical or used containers. Do not scatter bait on manure heaps or dispose of treated sacks/cardboard/fabric with manure. Clean up spillage promptly. Very highly toxic to bees.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

Disposal - 350 g Pack:

Shake container empty when completing use and dispose of empty container by wrapping with paper and putting in the garbage.

Disposal - 2 Kg pack:

Triple or preferably pressure rinse containers before disposal. Dispose of rinsings in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. If recycling replace cap and return clean containers to recycler or designated collection point. If not recycling break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit. Empty containers and product should not be burnt.

Treated sacks, cardboard or fabric may be re-used as necessary. After use they should be disposed of by wrapping in paper, placing in a plastic bag and putting in the garbage.

SAFETY DIRECTIONS

Will irritate eyes. Avoid contact with eyes. If product in eyes wash it out immediately with water. Wash hands after use. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and rubber gloves. After each days use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26; New Zealand 03 4747000).

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet

EXCLUSION OF LIABILITY

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

APVMA Approval No. 57892/0704

QuickBayt[®] is a Registered Trademark of Bayer.

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE 1800 033 111

DIRECTIONS FOR USE**Restraint**

DO NOT place bait in areas accessible to food producing animals or pets.

PEST	Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)
RATE	2.0 g/m ²
SITUATION	CRITICAL COMMENTS
<p>Outside: Abattoirs, animal housing, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around: Industrial sites, feedlot manure heaps, garbage dumps</p>	<p>Scatter application: The bait should be scattered (do not put in piles) onto dry level surfaces (eg. pathways, window ledges or in shallow dishes). Avoid extremely dusty areas.</p> <p>Bait Station application Set up bait stations by sprinkling into shallow tins, lids or similar containers and place where flies congregate. Position bait stations out of reach of animals or livestock and away from feed troughs, storage or mixing areas.</p> <p>Wet Bait: In many situations more flies can be attracted to moist bait. This can be achieved by sprinkling bait on moist hessian sacks (placed out of reach of all livestock) or by lightly moistening bait with a light spray of water.</p> <p>Paint-on application: To prepare a paste for painting, mix 50 g of QuickBayt with 37.5 mL of water. 15 minutes after mixing the paste can be applied with a brush to spots where flies rest (eg. Walls, joists, window ledges, stable fixtures). Additionally it can be painted onto hessian bags or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>Re-baiting: Re-baiting may be necessary in 3-5 days depending on exposure, weather and density of fly populations.</p>
<p>Inside: Commercial and industrial facilities and animal houses ONLY in areas inaccessible to livestock and domestic animals</p> <p>Under: Animal and poultry cages ONLY in areas inaccessible to livestock</p>	<p>Apply as a scatter bait, in bait stations or as a paint-on application as above.</p> <p>Place out of reach of all livestock and away from feed.</p> <p>Re-bait dry bait stations as above.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Main panel



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



QuickBayt[®] Spray

Fly Bait

ACTIVE CONSTITUENTS: 100 g/kg IMIDACLOPRID
1.0 g/kg (Z)-9-TRICOSENE

GROUP 4A INSECTICIDE

*** NET**

A water dispersible granule formulation to be mixed with water and used as a spray bait formulation for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTIONS FOR USE. Will visibly kill flies for 4 weeks.

For primary packs containing measuring packs (250 g (4x62.5 g); 1 kg (4x250 g)):

Contains 4 measuring packs which it is illegal to sell separately.

*12.5 g, 62.5 g, 250 g, 500 g, 750 g, 1 kg, 1.25 kg, 1.5 kg, 250 g (4x62.5 g), 1 kg (4x250 g)

Booklet / ancillary label

(Note: the rates shown in the Directions for Use table change with pack size and for this reason, there is a separate Direction for Use table for each pack size.)

DIRECTIONS FOR USE (12.5 g pack)**RESTRAINTS**

DO NOT apply in a manner which may cause contamination of animal feed.

DO NOT apply spray treatment outdoors if rain is expected within 48 hours.

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse).

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<p>Outside: Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around: Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside: Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	Nuisance flies including: Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)	12.5 g / 100 mL water (spray on application)	<p>Mix 12.5 g of QuickBayt® Spray with 100 mL of water and apply as a course spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 5 m² it is recommended that 100 mL of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor areas greater than 5 m² needs to be recalculated on the basis of 100 mL mixed bait per 5 m² floor area.</p> <p>Note: Around feedlot manure heaps DO NOT apply bait on manure heaps or dispose of treated targets with manure. See also Note on Application below.</p>
		12.5 g / 10 mL water (paint on application)	<p>To prepare a paste for painting, mix 12.5 g of QuickBayt® Spray with 10 mL of water. The mix can be applied with a brush to spots where flies rest (e.g. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 5 m² it is recommended that 10 mL of mix is applied in 3 to 5 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 5 m² need to be recalculated on the basis of 10 mL mixed bait per 5 m² floor area.</p> <p>See also Note on Application below.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

DIRECTIONS FOR USE (62.5 g, 250 g (4x62.5 g), 250 g, 500 g, 750 g, 1 kg, 1.25 kg and 1.5 kg packs)

RESTRAINTS

DO NOT apply in a manner which may cause contamination of animal feed.

DO NOT apply spray treatment outdoors if rain is expected within 48 hours.

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse).

SITUATION	PEST	RATE	CRITICAL COMMENTS
<p>This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.</p>			
<p>Outside: Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around: Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside: Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	<p>Nuisance flies including: Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)</p>	<p>62.5 g / 500 mL water (spray on application)</p>	<p>Mix 62.5 g of QuickBayt® Spray with 500 mL of water and apply as a course spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 25 m² it is recommended that 500 mL of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor area greater than 25 m² needs to be recalculated on the basis of 500 mL mixed bait per 25 m² floor area.</p> <p>Note: Around feedlot manure heaps DO NOT apply bait on manure heaps or dispose of treated targets with manure. See also Note on Application below.</p>
		<p>62.5 g / 50 mL water (paint on application)</p>	<p>To prepare a paste for painting, mix 62.5 g of QuickBayt® Spray with 50 mL of water. The mix can be applied with a brush to spots where flies rest (e.g. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 25 m² it is recommended that 50 mL of mix is applied in 12 to 25 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 25 m² need to be recalculated on the basis of 50 mL mixed bait per 25 m² floor area. See also Note on Application below.</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION.**

DIRECTIONS FOR USE (1 kg (4x250 g))**RESTRAINTS**

DO NOT apply in a manner which may cause contamination of animal feed.

DO NOT apply spray treatment outdoors if rain is expected within 48 hours.

DO NOT apply onto material which birds are likely to feed on directly (i.e. edible refuse).

SITUATION	PEST	RATE	CRITICAL COMMENTS
This product is not a conventional residual spray and therefore does not work by flies simply contacting a sprayed surface. The attractants within this bait mean that flies will be attracted to and subsequently attempt to feed on treated surfaces; they will then pick up a lethal dose. Because of this bait function, only portions of vertical surfaces need to be treated.			
<p>Outside: Abattoirs, animal housing, domestic dwellings, garbage containers, food processing plants and other industrial and commercial areas</p> <p>Around: Industrial sites, feedlot manure heaps, garbage dumps, poultry farms and animal housing (ONLY in areas inaccessible to livestock, domestic animals and children)</p> <p>Inside: Commercial and industrial facilities and animal houses (ONLY in areas inaccessible to livestock, domestic animals and children)</p>	Nuisance flies including: Adult houseflies (<i>Musca domestica</i>) and Lesser houseflies (<i>Fannia canicularis</i>)	250 g / 2 L water (spray on application)	<p>Mix 250 g of QuickBayt Spray with 2 L of water and apply as a course spray to areas where flies rest. Maintain agitation of suspension during application to ensure thorough mixing.</p> <p>For floor areas of 100 m² it is recommended that 2 L of mix is applied to approximately 1/3 of the surrounding (mainly vertical) surfaces. Any floor area greater than 100 m² needs to be recalculated on the basis of 2 L mixed bait per 100 m² floor area.</p> <p>Note: Around feedlot manure heaps DO NOT apply bait on manure heaps or dispose of treated targets with manure. See also Note on Application below.</p>
		250 g / 200 mL water (paint on application)	<p>To prepare a paste for painting, mix 250 g of QuickBayt Spray with 200 mL of water. The mix can be applied with a brush to spots where flies rest (e.g. walls, joists, window ledges, stable fixtures). Additionally it can be painted onto pieces of timber, or sheets of cardboard or fabric which are hung in areas frequented by flies. After use the brush can be cleaned with warm water.</p> <p>For floor areas of 100 m² it is recommended that 200 mL of mix is applied in 50 to 100 stripes to the surrounding (mainly vertical) surfaces. Any areas greater than 100 m² need to be recalculated on the basis of 200 mL mixed bait per 100 m² floor area. See also Note on Application below.</p>

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

(All packs)

GENERAL INSTRUCTIONS

QuickBayt® Spray can be applied in many locations. Avoid cool or windy locations because flies prefer warm sites for resting. For best results, time applications to begin at the start of the season before fly populations have reached their peak.

Fly control should be considered part of an integrated strategy which incorporates good sanitation, appropriate management of breeding sites (e.g. manure and other decaying organic matter) and exclusion measures for preventing access of flies into internal areas.

Note on Application:

Avoid applying this product to dusty/dirty surfaces. If the treated area becomes covered by dust/dirt, the product can be reactivated by gently spraying water mist on the treated surfaces. DO NOT store diluted insecticide. Prepare a fresh solution as necessary. DO NOT spray the moving parts of any machinery, electric motors or switchgear with the water-based product.

Re-baiting:

Re-apply bait when flies are no longer visibly affected in areas where the product has been applied (usually after about 4 to 6 weeks). For season-long control, monthly applications may be required, depending on the situation and duration of the fly season.

Cleaning Spray Equipment:

Clean spray equipment by rinsing thoroughly with lukewarm water. DO NOT allow washings to enter sewer drains or nearby water bodies.

WARNING

In some situations mould may develop (some time after application) on areas where this product has been used. If appearance of mould on a wall or fixture is a concern in the area of application then it is recommended that a surrogate target such as a disposable piece of timber, card or other surface is treated in its place.

A suitable detergent and cloth may also be used to remove mould in some areas. Dispose of cloth after use.

INSECTICIDE RESISTANCE WARNING

GROUP	4A	INSECTICIDE
-------	----	-------------

For insecticide resistance management, **QuickBayt** is a Group 4A insecticide. Some naturally occurring insect biotypes resistant to **QuickBayt** and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if **QuickBayt** or other Group 4A insecticides are used repeatedly. The effectiveness of **QuickBayt** on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Bayer CropScience Pty. Ltd. accepts no liability for any losses that may result from the failure of **QuickBayt** to control resistant insects. For further information on managing insecticide resistance contact your local supplier or Bayer Environmental Science or Bayer Animal Health representative.

PRECAUTIONS

DO NOT allow contact with raw or processed food, utensils or packaging.
DO NOT apply to surfaces within the reach of animals and children.
DO NOT spray directly on animals, feed or drinking supplies, or on milking equipment.
DO NOT apply past the point of run-off.

PROTECTION OF LIVESTOCK

DO NOT apply bait in locations accessible to domestic animals or livestock. Imidacloprid is dangerous to bees. **QuickBayt® Spray** will kill bees in hives which are over-sprayed or reached with spray drift.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT

DO NOT apply bait in locations accessible to birds. DO NOT contaminate waterways with the chemical or used containers. Clean up spillage promptly.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight.

(For foil bags and 12.5 g and 62.5 g bottle)

Single rinse before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Dispose of empty containers by wrapping with paper and putting in garbage. Dispose of any rinsates or any undiluted chemical according to State/Territory legislative requirements. Empty container and product should not be burnt. DO NOT re-use empty container for any other purpose.

(For bottles >62.5 g)

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

If washing of treated surfaces is necessary, any wash-off water should be prevented from entering waterways or drains. Wash water should be collected and disposed as recommended above.

SAFETY DIRECTIONS

When opening the container, preparing and using the prepared spray or paste wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length chemical resistant gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet which may be found at www.bayeres.com.au.

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty. Ltd. accepts no liability or responsibility for loss or damage from failure to follow such directions and instructions.

APVMA Approval No.: 61737/58686

QuickBayt® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6800
Technical enquiries: 1800 804 479
Website: www.bayer.com.au

For 24 hour specialist advice in emergency only phone 1800 033 111

Batch Number:

DOM:

BARCODE

Main panel of measuring packs

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



QuickBayt[®] Spray

Fly Bait

**ACTIVE CONSTITUENTS: 100 g/kg IMIDACLOPRID
1.0 g/kg (Z)-9-TRICOSENE**

GROUP 4A INSECTICIDE

*** g NET**

A water dispersible granule formulation to be mixed with water and used as a spray bait formulation for the control of adult houseflies and lesser houseflies (including organophosphate resistant strains) in commercial, industrial and domestic areas, as per the DIRECTIONS FOR USE. Will visibly kill flies for 4 weeks.

**NOT TO BE SOLD SEPARATELY.
BEFORE USE READ ALL DIRECTIONS ON THE OUTER PACK.**

***62.5 g, 250 g**

Back panel of measuring packs**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well ventilated area out of direct sunlight. Single rinse before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. Dispose of empty containers by wrapping with paper and putting in garbage. Dispose of any rinsates or any undiluted chemical according to State/Territory legislative requirements. Empty container and product should not be burnt. DO NOT re-use empty container for any other purpose.

If washing of treated surfaces is necessary, any wash-off water should be prevented from entering waterways or drains. Wash water should be collected and disposed as recommended above.

SAFETY DIRECTIONS

When opening the container, preparing and using the prepared spray or paste wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), a washable hat and elbow-length chemical resistant gloves. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

APVMA Approval No.: 61737/58686

QuickBayt® is a Registered Trademark of the Bayer Group.

Bayer CropScience Pty. Ltd.
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn, Vic. 3123
Phone: (03) 9248 6888
Fax: (03) 9248 6803

Distributed by:
Bayer Animal Health
Technical enquiries: 1800 678 368
Website: www.bayeranimal.com.au

For 24 hour specialist advice in emergency only phone 1800 033 111